

**Heartsaver AED  
Child CPR and Child AED Section**

21. You find an unresponsive child. You open the airway and find that the child is not breathing. You try to give rescue breaths, but the chest does not rise. What should you do?
- give several cycles of chest compressions
  - re-open the airway and try the breath again
  - attach an AED
  - look, listen, and feel for normal breathing
22. How often should you give rescue breaths without chest compressions for a child who is not breathing but has other signs of circulation?
- about once every second
  - about once every 3 seconds
  - about once every 6 seconds
  - about once every 10 seconds
23. Which of the following best describes good rescue breaths for a child?
- each breath is fast and you see the chest rise
  - each breath is slow and you see the stomach rise
  - each breath is slow and you see the chest rise
  - each breath is fast and you see the stomach rise
24. A 7-year-old child is pulled from below the surface of a neighbor's pool. The child does not respond. You send someone to phone 911. What should you do next?
- turn the child's head down and do the Heimlich maneuver to drain water from the lungs
  - give 5 back blows and 5 chest thrusts; then do a finger sweep of the mouth
  - using proper hand position give chest compressions at a rate of at least 100 times per minute
  - open the airway, check breathing, and if she is not breathing normally, give 2 breaths
25. A 4-year-old child is found unresponsive and not breathing, so you give 2 good rescue breaths and check for signs of circulation. What are the signs of circulation?
- response to shouting
  - normal breathing, coughing, or movement
  - response to chest compression
  - bluish lips or skin



26. You think a 5-year-old child is choking. Her lips are turning blue, and she cannot speak when you ask questions. She is holding her neck and nods when you ask if she is choking. You tell her you will help. What should you do next?
- give several cycles of 5 back blows and then 5 chest thrusts
  - use your finger to sweep out the child's mouth; then give rescue breaths
  - do nothing until the child stops breathing and goes limp in your arms
  - give abdominal thrusts until the object pops out or she becomes unresponsive
27. When should you phone 911 if you are alone and you find an unresponsive infant or child?
- phone 911 immediately
  - phone 911 after you give CPR for about 5 minutes
  - phone 911 before you start CPR
  - phone 911 after you give CPR for about 1 minute
28. Where should you press when you give a child chest compressions during CPR?
- over the lower half of the breastbone
  - over the belly button
  - over the top half of the breastbone
  - between the belly button and the bottom of the breastbone
29. The AED case at your workplace contains both adult and child AED pads. Which of the following lists the correct ages for use of the child AED pads?
- use child AED pads for a victim any age
  - use child AED pads for a victim from age 1 up to 8 years of age
  - use child AED pads for any victim 8 years of age or older
  - you should not use child AED pads with any AED

**END OF CHILD SECTION**